Turiva®





CLINICAL APPLICATIONS

- Maintains Normal Inflammatory Balance
- Supports Microbiome Diversity
- · Potent Antioxidant Action
- Enhances and Balances Immune Function



IMMUNE HEALTH

Turiva® contains an exclusive blend of bioactive phytonutrients that make up the Complete Turmeric Matrix for increased clinical efficacy. It is a natural complex that includes curcumin, turmerones, turmerin, curcumol, cyclocurcumin, curdione and B-elemene making it a comprehensive, immune-boosting turmeric supplement.

Overview

Modern lifestyle choices, such as an unbalanced diet and poor nutrition, have a major impact on health determinants. Immune and inflammatory challenges arise when our bodies cannot keep up with the demands of handling stress, toxin removal and cellular function preservation. Turiva® provides a well-rounded blend of turmeric phytonutrients that can help support normal inflammatory input and strengthen immune reserves.¹ The formula targets various proinflammatory markers and supports healthy breathing, cognitive function, gut and immune function.¹,² The molecular structure of turmeric's components–proteins, fibers, carbohydrates and volatile oils–contributes to its bioefficacy and maximizes its pleiotropic effects, making it an essential phytonutrient to combat various inflammatory pathways in our body.

Turmeric (Complete Turmeric Matrix)†


Whole-root turmeric and its active components have been used in traditional Ayurvedic medicine for centuries. In herbal medicine of old, practitioners used teas, tinctures and extracts of all types. In the 21st century, as research grew on the benefits of turmeric, the focus shifted to identifying and isolating one individual compound, curcumin, rather than delivering the comprehensive benefits of a matrix of turmeric bioactives. As a result, concentrating curcumin led to poor absorption, and

pharmaceutical methods were applied to bypass the gut and increase its bioavailability. The glaring disadvantage of applying this pharmaceutical model to botanicals is that it misses the benefits of other bioactives present within the turmeric matrix and their positive effects on the microbiome. New research on turmeric shows the additional bioactives in turmeric have additional benefits and enhance bioavailability. The **Complete Turmeric Matrix** includes compounds from the entire turmeric root, all working together as nature intended to deliver better results.

- **Curcumin:** Activates alkaline phosphatase that decreases LPS activity and inhibits the expression of the NF-kB/lkB pathway.^{3,4}
- **Turmerones:** Enhance the bioavailability and bioactivity of curcumin ⁵
- **Turmerin:** Inhibits oxidative DNA damage, formation of lipid peroxides, and arachidonic acid release.⁶
- **Curcumol:** Inhibits oxidative stress and targets key signaling pathways MAPK/ERK, PI3K/Akt and NF-kB.⁷
- **Cyclocurcumin:** Exhibits synergistic bioactivity with curcumin and inhibits the release of TNF-alpha.^{8,9}
- **Curdione:** Inhibits the production of prostaglandin E2 (PGE2) and COX-2 expression.¹⁰
- **B-elemene:** Downregulates pro-inflammatory cytokines TNF-alpha, IFN, TGF-beta, IL-6 and IL-10.^{11,12}

This matrix of bioactive compounds supports a healthy GI tract, enhances detoxification, creates a healthy microbiome, and helps maintain normal inflammatory balance.





Turmeric and Immune Health[†]


Turmeric and its phytonutrients, like curcumin, have been shown to support lowering of lipid peroxidation by maintaining the activity of antioxidant enzymes, such as superoxide dismutase, catalase and glutathione peroxidase, at higher levels. In animal studies, curcumin was reported to decrease the release of reactive oxygen species from macrophages and inhibit histamine release from mast cells. The chemical structure of the curcumin molecule allows it to have a significant effect on oxidative stress and immune-mediated reactions. However, several findings indicate that antioxidant and protective effects are synergistically enhanced when other turmeric components are included. Other turmeric components, such as turmerones, curdione, curcumol, cyclocurcumin, such as turmerones, curdione, curcumol, cyclocurcumin, such as turmerones, such and β -elemene, li,12,25 all have significant impact on immune health and maintaining normal inflammatory balance.

A complete turmeric matrix was tested for its impact on immune cells. The test results showed it had a three-fold increase in phagocytic ability.¹⁷ Due to its pleiotropic effects and ability to pursue multiple inflammatory targets, turmeric has proven uses as an immune-boosting agent. A meta-analysis of eight randomized, controlled trials indicated that doses of 1,000 mg of curcumin per day were effective in balancing overactive immune challenges.¹⁹ In a randomized, double-blind, placebo-controlled clinical trial published in the Journal of Medicinal Food, a total of 36 patients with immune challenges were included to test out the efficacy of a complete turmeric matrix. A dose of 250 mg or 500 mg of the turmeric product was given twice a day to the patients in the treatment groups. Specific objective measures and lab markers were used to assess the response of these patients. The results indicated that a complete turmeric formulation improved objective measurements and maintained normal inflammatory balance.¹⁸ (See Figure: 1)

Turmeric and Gut Health[†]

All sources of orally consumed nutrients must pass through the gut before moving along to the rest of the body. Likewise, the gut microbiome is now considered influential in every aspect of human metabolism, making it an important mediator of phytonutrients. Therefore, the gut and its microbiota are the primary targets of medicinal compounds like turmeric. Turmeric and its phytonutrients, like curcumin, have been shown to support expression of protective proteins such as Nrf2. They also support normal balance of the expression of various inflammatory agents in the gut, such as NFkB, COX enzyme(s), and matrix metalloproteinase-9. In addition, they boost NK cell activity. The mechanism of action of curcumin is that it can indirectly bind to toll-like receptor 4 (TLR4). Literature indicates that this action supports a balance in the innate immune response to

Immune Modulation with CTM





lipopolysaccharides (LPS).²⁰ Furthermore, turmeric can help attenuate LPS-induced cytokine release and tight junction protein disruption. Other, non-curcuminoid compounds, such as curdione and germacrone, also help balance LPS-induced immune challenges.²² This promotes overall gut mucosal health and helps create equilibrium in inflammatory responses.⁴

Turmeric Bioactivity vs. Bioefficacy[†]

Bioavailability is defined as the delivery of a biologically active compound to a target tissue; it is not simply the absorption of higher amounts of the nutrient into the blood. There has been continuous debate on the bioavailability of curcumin, and it is important to note that most clinical research of turmeric-derived compounds is focused on curcuminoid activity.¹³ Curcuminoids undergo extensive metabolism during and after ingestion.¹³ This is one of the key reasons curcumin is encouraged to be consumed with an adjuvant or made with "bioavailability enhancers" that dilute the breakdown process. However, these enhanced versions show no significant impact in human clinical studies. Studies showing increased absorption have not demonstrated increased positive clinical outcomes. Based on this, it is important to shift focus to bioefficacy, or the effectiveness of a material in a biological environment, over bioavailability. Components of the whole turmeric root have proven that the therapeutic effect of turmeric extends beyond curcuminoids alone.16

Directions

1 capsule per day or as recommended by your health care professional.

Does Not Contain

Gluten, corn, yeast, artificial colors or flavors.

Caution

Do not consume this product if you are pregnant or nursing.

Supplement Facts Serving Size 1 Capsule Servings Per Container 60 & 120 Amount Per Serving Value Turmeric Root Extract 500 mg (Complete Turmeric Matrix) (Standardized to contain 45-55% Curcuminoids, 3-8% Volatile Oil, 2-6% Turmerin) * Daily Value not established.

Other Ingredients: Hypromellose (Natural Vegetable Capsule), Microcrystalline Cellulose, Magnesium Stearate and Silicon Dioxide.

ID# 172060 60 Capsules ID# 172120 120 Capsules

References

- Aggarwal BB, Harikumar KB. Potential therapeutic effects of curcumin, the anti-inflammatory agent, against neurodegenerative, cardiovascular, pulmonary, metabolic, autoimmune and neoplastic diseases. *The International Journal of Biochemistry & Cell Biology*. 2009;41(1):40-59. doi:https://doi.org/10.1016/j.biocel.2008.06.010
- 2. Suzuki M, Nakamura T, Iyoki S, et al. Elucidation of Anti-allergic Activities of Curcumin-Related Compounds with a Special Reference to Their Anti-oxidative Activities. *Biological & Pharmaceutical Bulletin*. 2005;28(8):1438-1443. doi:https://doi.org/10.1248/bpb.28.1438
- Ghosh SS, Bie J, Wang J, Ghosh S. Oral supplementation with nonabsorbable antibiotics or curcumin attenuates western diet-induced atherosclerosis and glucose intolerance in LDLR-/- mice--role of intestinal permeability and macrophage activation. PLoS One. 2014 Sep 24;9(9):e108577. doi: 10.1371/journal.pone.0108577. PMID: 25251395; PMCID: PMC4177397.
- Wang Y, Tang Q, Duan P, Yang L. Curcumin as a therapeutic agent for blocking NF-κB activation in ulcerative colitis. Immunopharmacol Immunotoxicol. 2018 Dec;40(6):476-482. doi: 10.1080/08923973.2018.1469145. Epub 2018 Aug 15. PMID: 30111198.
- 5. Cao W, Chen X, Xiao C, Lin D, Li Y, Luo S, Zeng Z, Sun B, Lei S. Arturmerone inhibits the proliferation and mobility of glioma by downregulating cathepsin B. *Aging* (Albany NY). 2023 Sep 26;15(18):9377-9390. doi: 10.18632/aging.204940. Epub 2023 Sep 26. PMID: 37768200; PMCID: PMC10564430.
- Srinivas L, Shalini VK, Shylaja M. Turmerin: a water soluble antioxidant peptide from turmeric [Curcuma longa]. Arch Biochem Biophys. 1992 Feb 1;292(2):617-23. doi: 10.1016/0003-9861(92)90040-4. PMID: 1731625.
- 7. Wei W, Rasul A, Sadiqa A, Sarfraz I, Hussain G, Nageen B, Liu X, Watanabe N, Selamoglu Z, Ali M, Li X, Li J. Curcumol: From Plant Roots to Cancer Roots. *Int J Biol Sci.* 2019 Jun 4;15(8):1600-1609. doi: 10.7150/ijbs.34716. PMID: 31360103; PMCID: PMC6643219.
- 8. Kiuchi F, Goto Y, Sugimoto N, Akao N, Kondo K, Tsuda Y. Nematocidal activity of turmeric: synergistic action of curcuminoids. *Chem Pharm Bull* (Tokyo). 1993 Sep;41(9):1640-3. doi: 10.1248/cpb.41.1640. PMID: 8221978.
- 9. Fu M, Chen L, Zhang L, Yu X, Yang Q. Cyclocurcumin, a curcumin derivative, exhibits immune-modulating ability and is a potential compound for the treatment of rheumatoid arthritis as predicted by the MM-PBSA method. *Int J Mol Med.* 2017 May;39(5):1164-1172.
- Oh OJ, Min HY, Lee SK. Inhibition of inducible prostaglandin E2 production and cyclooxygenase-2 expression by curdione from Curcuma zedoaria. *Arch Pharm Res.* 2007 Oct;30(10):1236-9. doi: 10.1007/BF02980264. PMID: 18038902.
- 11. Tan T, Liu J, Luo R, Wang R, Yin L, Liu M, Zeng Y, Zeng Z, Xie T. Recent Advances in Understanding the Mechanisms of Elemene in Reversing Drug Resistance in Tumor Cells: A Review. *Molecules*. 2021 Sep 24;26(19):5792. doi: 10.3390/molecules26195792. PMID: 34641334; PMCID: PMC8510449.



- Nair A, Amalraj A, Jacob J, Kunnumakkara AB, Gopi S. Non-Curcuminoids from Turmeric and Their Potential in Cancer Therapy and Anticancer Drug Delivery Formulations. *Biomolecules*. 2019 Jan 2;9(1):13. doi: 10.3390/biom9010013. PMID: 30609771; PMCID: PMC6358877.
- 13. Heger M, van Golen RF, Broekgaarden M, Michel MC. The Molecular Basis for the Pharmacokinetics and Pharmacodynamics of Curcumin and Its Metabolites in Relation to Cancer. *Pharmacological Reviews*. 2013;66(1):222-307. doi:https://doi.org/10.1124/pr.110.004044
- Takeda Y, Bhardwaj A. Nonsteroidal Anti-Inflammatory Agents Differ in Their Ability to Suppress NF-kappaB Activation, Inhibition of Expression of cyclooxygenase-2 and Cyclin D1, and Abrogation of Tumor Cell Proliferation. Oncogene. Published December 9, 2004. https://pubmed.ncbi.nlm.nih.gov/15489888/
- Swarnakar S, Ganguly K, Kundu P, Banerjee A, Maity P, Sharma AV. Curcumin Regulates Expression and Activity of Matrix Metalloproteinases 9 and 2 during Prevention and Healing of Indomethacin-induced Gastric Ulcer. *Journal of Biological Chemistry*. 2004;280(10):9409-9415. doi:https://doi.org/10.1074/jbc. m413398200
- Gopi S, Jacob J, Varma K, et al. Comparative Oral Absorption of Curcumin in a Natural Turmeric Matrix with Two Other Curcumin Formulations: An Open-label Parallel-arm Study. *Phytotherapy Research*. 2017;31(12):1883-1891. doi:https://doi.org/10.1002/ ptr.5931
- 17. Gopi, S, et al. (2014). Cell culture study on the cytotoxic effects of "Cureit"- A novel bio available curcumin-anti cancer effects. 6. 96-100.
- Amalraj A, Varma K, Jacob J, et al. A Novel Highly Bioavailable Curcumin Formulation Improves Symptoms and Diagnostic Indicators in Rheumatoid Arthritis Patients: A Randomized, Double-Blind, Placebo-Controlled, Two-Dose, Three-Arm, and Parallel-Group Study. *Journal of Medicinal Food*. 2017;20(10):1022-1030. doi:https://doi.org/10.1089/jmf.2017.3930
- 19. Daily JW, Yang M, Park S. Efficacy of Turmeric Extracts and Curcumin for Alleviating the Symptoms of Joint Arthritis: A Systematic Review and Meta-Analysis of Randomized Clinical Trials. *Journal of Medicinal Food*. 2016;19(8):717-729. doi:https://doi.org/10.1089/jmf.2016.3705
- 20. Burge K, Gunasekaran A, Eckert J, Chaaban H. Curcumin and Intestinal Inflammatory Diseases: Molecular Mechanisms of Protection. *International Journal of Molecular Sciences*. 2019;20(8):1912. doi:https://doi.org/10.3390/ijms20081912
- 21. Sandur SK, Pandey MK, Sung B, et al. Curcumin, demethoxycurcumin, bisdemethoxycurcumin, tetrahydrocurcumin and turmerones differentially regulate anti-inflammatory and anti-proliferative responses through a ROS-independent mechanism. *Carcinogenesis*. 2007;28(8):1765-1773
- 22. Oh OJ, Min HY, Lee SK. Inhibition of inducible prostaglandin E2 production and cyclooxygenase-2 expression by curdione from Curcuma zedoaria. *Arch Pharm Res.* 2007;30(10):1236-1239

- 23. Xia Q, Wang X, Xu DJ, Chen XH, Chen FH. Inhibition of platelet aggregation by curdione from Curcuma wenyujin essential Oil. *Thromb Res.* 2012;130(3):409-414
- 24. Lekshmi PC, Arimboor R, Raghu KG, Menon AN. Turmerin, the antioxidant protein from turmeric (Curcuma longa) exhibits antihyperglycaemic effects. *Nat Prod Res.* 2012;26(17):1654-1658.
- 25. Liu J, Zhang Z, Gao J, Xie J, Yang L, Hu S. Downregulation effects of beta-elemene on the levels of plasma endotoxin, serum TNF-alpha, and hepatic CD14 expression in rats with liver fibrosis. *Front Med.* 2011;5(1):101-105.

LT-PDN-030-F

